

THE SPECIFIC SPECTRAL DATA OF DOMINANT TREES IN PEAT-FOREST IN CENTRAL KALIMANTAN, INDONESIA

Hendrik Segah¹, Hiroshi Tani², Muhammad Evri³, Aswin Usup⁴, Kazuyo Hirose¹, and Mitsuru Osaki^{1,2}

¹) Center for Sustainability Science (CENSUS), Hokkaido University, Kita-8, Nishi-9, Kita-ku, Sapporo 060-0809 Japan, Phone: +81-11-706-E-mail: segah@census.hokudai.ac.jp

²) Graduate School of Agriculture, Hokkaido University, Japan

³) Agency for the Assessment and Application of Technology (BPPT), Indonesia

⁴) Faculty of Agriculture, University of Palangka Raya (UNPAR), Indonesia

SUMMARY

This paper reports results of the compilation of the spectral library of dominant trees in peat-forest in Central Kalimantan, Indonesia supported under JST-JICA Project: “*Wild Fire and Carbon Management in Peat Forest in Indonesia*”. Spectral libraries are commonly established as a means to archive representative signatures of natural materials. Such signatures can then be used to train feature extraction and classification algorithms applied to remote sensing data imagery. Spectral analyses were performed to the 33 dominant species in peat-forest in Central Kalimantan, Indonesia obtained during the field survey measurement (in-situ measurements) in 2011. We found, each vegetation type has unique spectral signature or characteristics. The shapes of reflectance spectra can be used for identification of vegetation types combine with remote sensing data.

KEY WORDS Spectral library/signature, Peat-Forest, Spectroradiometer, Vegetation, Central Kalimantan

INTRODUCTION

Remote sensing applications in precision identification of agriculture and forest vegetation have been steadily increasing in recent years due to improvements in spatial and spectral resolutions of remotely sensed imagery. There is a rapidly growing interest in methods for automatic plant identification, especially in agricultural and forestry research. Innovations in sensors are permitting the connection of remote sensing with methods of laboratory spectroscopy (Raymond et al., 2003; Kuusk et al., 2004). The Spectral library is “a set of class or end member spectrum to be referred to during feature extraction process from hyperspectral data”. It can be further explained as a collection of spectral plots which is being compiled so that it can be used as a reference in information extraction from hyperspectral data. The reflection of radiation from the forest understorey and ground vegetation, in particular, plays a significant role in the forming of total forest reflectance (Danson & Curran, 1993).

Field campaign is the most common and easiest way to collect object spectra of different land cover types or vegetation, and this method produces results which can be easily compiled into a spectral library (Kuusk et al., 2004; Lau & Hashim, 2007). To carry out a field campaign, a

spectroradiometer is needed to record the spectra received from the objects found, especially tropical forest vegetation (tree species), in the study area.

MATERIAL AND METHODS

Study Area

The study area is located in a tropical peat swamp forest area near Palangka Raya, the capital of Central Kalimantan Province, Indonesia (Fig. 1) and approximately 120 km from the Java sea with an elevation between 15-20 m above sea level (Shimada et al., 2001; Weiss et al., 2002). Peat swamps cover extensive areas of lowland Kalimantan, with estimates varying between 8% and 11% (McKinnon et al., 1996). Area of Central Kalimantan is 153,564 km², comprising mostly jungle (ca. 126,200 km², 82.18%), swamps (11.80%), rivers and lakes (2.97%), and agricultural land (3.05%). Central Kalimantan contains about three million hectares of peatland which is one of the largest unbroken peatland areas in the world and is located between 0° 45' N and 3° 3' S, between 111° and 116° E. Ground surveys and field investigations were carried out in the study area in July 2011.




Figure 1. Map of study area in Central Kalimantan Province, Indonesia with an extent of 75 km x 72 km or approximately 540,000 ha.

Methods

The measurements of the reflectance spectra of ground vegetation were performed by an Analytical Spectral Devices (ASD) FieldSpec Pro-FR Spectroradiometer (FieldSpec 3 FR)

with a spectral range of 350-2500 nm and a rapid data collection time of 0.1 second per spectrum. The main purpose of this study is to develop the spectral library of individual tree of peat-forest using a ground-based spectroradiometer. The instrument uses three detectors spanning the visible and near infrared (VNIR) and shortwave infrared (SWIR1 and SWIR2). The FieldSpec spectroradiometer is specifically designed for field environment remote sensing to acquire visible near-infrared (VNIR) and short-wave infrared (SWIR) spectra.



Figure 2. Spectral library data collection activities (field survey and sample measurements)

The sensor, with a field of view of 25°, was positioned 30-40 cm above the samples at nadir position. Prior to each three measurements, a white reference panel with approximately 100% reflectance was used as a reference standard. The measurements were conducted under clear and cloudless sky between 10:00 and 14:00 at local time (West Indonesian Time). A contact probe fore optic was used to minimise BRDF effects from shiny surfaces. Measurements were made with a standard laboratory set up (Pfitzner et al., 2006) using the high intensity contact probe at 30° of nadir. The ASD contact probe used is equipped with an internal light source (a halogen bulb light source 6.5 W, with colour temperature $2901 \pm 10\%$ K). Spot size of the Contact Probe has a diameter of 22 mm as shown in Figure 2.

The contact probe is an alternative method for collection of laboratory spectra because SWIR spectra collected are found to exhibit improved signal to noise (S:N) ratios, lower standard deviation and less intrinsic uncertainty than spectra collected using more traditional laboratory techniques (MacArthur, 2007). The spectrometer and contact probe were warmed prior to use for 90 minutes and 30 minutes, respectively. A white reference measurement was undertaken prior to each material (tree leaves) being sampled. Spectral measurements were made using 30 averages.

RESULTS

The samples of the library spectra measurements from 10 dominant trees species in peat-forest in Central Kalimantan are shown in Figure 3 from the total of 33 species during field survey and measurements in full-range wavelength (350 – 2500 nm). The definition of dominant trees are trees (or shrubs) with crowns receiving full light from above and partly from the side; usually larger than the average trees or shrubs in the stand, with crowns that extend above the general level of the canopy and that are well developed but possibly somewhat crowded on the sides. A dominant tree is one which generally stands head and shoulders above all other trees in its vicinity. However, there may be young, vigorous tree nearby, but not overtopped by a dominant tree. These smaller trees may be considerably shorter than the dominants, but still be receiving full light from above and partly from the sides. In its own immediate environment, it is dominant and should be recorded as such. Only understorey trees immediately adjacent to the overstorey tree will be assigned subordinate crown classes (Raulier et al., 2003; Miles et al., 2003).




Figure 3. Spectral reflectance's from 10 species of peat-forest in full-range wavelength (350 – 2500 nm)




Figure 4. Spectral reflectance's from 10 species of peat-forest in SWIR wavelength (1000 – 1800 nm)

The spectra demonstrate the basic varieties in the spectral shapes across the visible to shortwave infrared (350–2500 nm) wavelength ranges. These are examples from 33 species measured from various locations in the study areas. Figure 4 also presents examples of spectral reflectance from 10 dominant tree species of peat-forest in SWIR wavelength (1000–1800 nm) with special characteristic of Tumeh (*Combretocarpus rotundatus*) which was always in lower reflectance values than other tree species such as Belangeran (*Shorea belangeran*), Gerunggang (*Cratoxylon* sp.), Jelutung (*Dyera costulata*), Kayu Hitam (*Dyospiros* sp.), Manggis Hutan (*Garcinia* sp.), Ramin (*Gonystylus bancanus*), Mahang (*Macaranga* sp.), Nyatoh (*Palaquium* sp.), and Kayu Pelawan (*Tristania maingayi*).

As shown in Fig. 3 and Fig. 4, the variability is caused mainly by the spatial in-homogeneities of canopies and species. If we made a series of measurements without moving the spectrometer, the variability (caused by noise and possible changes in illumination) was several times less (Kuusk et al., 2004; Abbasia et al., 2008). However, even though very important, leaf optical properties alone are not adequate to explicitly spectrally distinguish tree species such as in the peat-forest ecosystem of Central Kalimantan. Also, other canopy components such as bark branches, twigs and understory must be considered.

DISCUSSION AND CONCLUSIONS

Spectral analysis and measurements were performed during the field campaign to 33 vegetation species spectra and found that the vegetation spectra are very sensitive to environmental parameters such as leaf condition, vigorous and other physiological and

biochemical parameters. Spectral libraries are data archives that consist of spectral signatures measured on selected natural and/or man-made materials. These libraries should also include the corresponding meta-information on the records. There are in principle two potential purposes: the use of the data in remote sensing as in-situ radiance/reflectance measurements for the calibration and/or end-members selection for further data processing (Selige *et al.* 2006).

The knowledge gained from laboratory and field studies of vegetation spectra and laboratory spectral analysis methods are directly applicable to remote sensing data. This analysis highlights the possible integration of in-situ hyperspectral measurements with airborne/space-borne hyperspectral remote sensing data for automatic identification and discrimination of various species/vegetation on the study areas. The continually updated spectral library of peatland forest provides initial collections of spectra covering the wavelength range 350–2500 nm that allows for the modeling of basic vegetation and land-cover properties by statistical inference. This study has produced the first preliminary spectral library of the dominant forest tree species of the peat-forest in Central Kalimantan (Indonesia) taking into account the range of spectral variability expected for the species measured under natural illumination conditions.

Meanwhile, spectral libraries from ground measurements and hyperspectral sensors can collect continuous spectral reflectance data that enable classification of peat-forest types, discrimination of plant/tree species (through biophysical parameters), forest degradation, soil types and so on could support the integrated Measurement, Reporting and Verification (MRV) of Reducing Emissions from Deforestation and Forest Degradation (REDD) programmes in Indonesia. Monitoring systems that allow for credible measurement, reporting and verification of REDD+ activities are among the most critical elements for the successful implementation of any REDD+ mechanism.

ACKNOWLEDGEMENTS

This research was supported by Hokkaido University, JST-JICA SATREPS Project “Wild Fire and Carbon Management in Peat-Forest in Indonesia” collaborate with the University of Palangka Raya (UNPAR) and Indonesia Agency for the Assessment and Application of Technology (BPPT). The authors would like to thank to Dr. Suwido H. Limin, Mr. Ici P. Kulu, Mr. Kitso Kusin and the staffs from the Centre for International Management of Tropical Peatland (CIMTROP) University of Palangka Raya, Central Kalimantan, Indonesia for their field support and assistance during the field campaign.

REFERENCES

- Abbasia, M., Schaepmanb, M.E., Darvishsefata, A., Bartholomeusb, H.M., Marvi Mohajera, M.R. and Sobhania, H. (2008) Spectroradiometric measurements of tree species in the Caspian Forests of Iran. Proceedings of the International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences. Vol. XXXVII. Part B7. Beijing, China.
- Lau, A. and Hashim, M. (2007) The Design and Building of Spectral Library of Tropical Rain Forest in Malaysia. Proceedings of the 28th Asian Conference on Remote Sensing, Kuala Lumpur, Malaysia, 12-16 November 2007. Asian Association of Remote Sensing (AARS).

Kuusik, A., Lang, M. and Nilson, T. (2004) Simulation of the reflectance of ground vegetation in sub-boreal forests. *Agricultural and Forest Meteorology* 126, 33–46.

Danson, F.M., and Curran, P.J. (1993). Factors affecting the remotely sensed response of coniferous forest plantations. *Remote Sensing of Environment* 43 (1), 55–65.

Raulier, F., Lambert, M-C., Pothier, D. and Ung, C-H. (2003) Impact of dominant tree dynamics on site index curves. *Forest Ecology and Management* 184, 65–78.

MacArthur, A. (2007). Field Guide for the ASD FieldSpec Pro-White Reference Mode. Version 2. Natural Environment Research Council Field Spectroscopy Facility. Available online at http://fsf.nerc.ac.uk/resources/guides/pdf_guides/asd_guide_v2_wr.pdf, (accessed on 1 March 2012)

Miles, R., Silman, J., Terborgh, W. and Kiltie, R.A. (2003) Population regulation of a dominant rain forest tree by a major seed predator. *Ecology*, 84 (2), 431–438.

Pfizer, K., Bollhöfer, A. and Carr, G. (2006). A standard design for collecting vegetation reference spectra: Implementation and implications for data sharing. *Journal of Spatial Sciences*, 52, 79–92.

Raymond, F., Kokaly, D., Despain, G., Clark, R.N. and Livo, E. (2003) Mapping vegetation in Yellowstone National Park using spectral feature analysis of AVIRIS data. *Remote Sensing of Environment* 84, 437–456.

Selige, T., Bohner, J. and Schmidhalter, U. (2006) High resolution topsoil mapping using hyperspectral image and field data in multivariate regression modeling procedures. *Geoderma*, 136, 235–244.

Shimada, S., Takahashi, H., Haraguchi, A. and Kaneko, M. (2001) The carbon content characteristics of tropical peats in Central Kalimantan, Indonesia; Estimating their spatial variability in density. *Biogeochemistry*, 53, 249–267.

Weiss, D., Shotyk, W., Rieley, J.O., Page, S., Gloor, M., Reese, S. and Martinez-Cortizas, A. (2002) The geochemistry of major and selected trace elements in a forested peat bog, Kalimantan, SE Asia, and its implications for past atmospheric dust deposition. *Geochimica et Cosmochimica Acta*, 66, 2307–2323.